

HADH2 Blocking Peptide

| Catalog # | Source | Reactivity | Applications |
|---------------------------|---|------------------|--------------|
| CBP1525 | Synthetic | H, M, R, B, D, P | BL |
| Description | The peptide is used to block Anti-HADH2 Antibody (#CPA1525) reactivity. | | |
| Form | Lyophilized powder | | |
| Gene Symbol | HSD17B10 | | |
| Alternative Names | ERAB; HADH2; MRPP2; SCHAD; XH98G2; 3-hydroxyacyl-CoA dehydrogenase type-2; 17-beta-hydroxysteroid dehydrogenase 10; 17-beta-HSD 10; 3-hydroxy-2-methylbutyryl-CoA dehydrogenase; 3-hydroxyacyl-CoA dehydrogenase type II; Endoplasmic reticulum-associated amyloid beta-peptide-binding protein; Mitochondrial ribonuclease P protein 2; Mitochondrial RNase P protein 2; Short-chain type dehydrogenase/reductase XH98G2; Type II HADH | | |
| Entrez Gene | 3028 (Human); 63864 (Rat) | | |
| SwissProt | Q92851 (Human) | | |
| Purity | >85% | | |
| Quality Control | The quality of the peptide was evaluated by reversed-phase HPLC and by mass spectrometry. | | |
| Directions for Use | Blocking Peptide to the diluted primary antibody in a molar ratio of 10:1 (peptide to antibody) and incubate the mixture at 4°C for overnight or at room temperature for 2 hours. | | |
| Storage/Stability | Shipped at 4°C. Store at -20°C for one year. | | |

Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC- Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference

Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb- Rabbit, S- Sheep, Z- Zebrafish

COHESION BIOSCIENCES LIMITED

WEB
www.cohesionbio.com

ORDER
order@cohesionbio.com

SUPPORT
techsupport@cohesionbio.com

CUSTOM
custom@cohesionbio.com