

CD184 (pS339) Blocking Peptide

| Catalog # | Source | Reactivity | Applications |
|---------------------------|--|----------------------|--------------|
| CBP2254 | Synthetic | H, M, R, B, D, Mk, P | BL |
| Description | The peptide is used to block Anti-CD184 (pS339) Antibody (#CPA2254) reactivity. | | |
| Form | Lyophilized powder | | |
| Gene Symbol | CXCR4 | | |
| Alternative Names | C-X-C chemokine receptor type 4; CXC-R4; CXCR-4; FB22; Fusin; HM89; LCR1; Leukocyte-derived seven transmembrane domain receptor; LESTR; Lipopolysaccharide-associated protein 3; LAP-3; LPS-associated protein 3; NPYRL; Stromal cell-derived factor 1 receptor; SDF-1 receptor; CD184 | | |
| Entrez Gene | 7852 (Human); 12767 (Mouse) | | |
| SwissProt | P61073 (Human); P70658 (Mouse); O08565 (Rat) | | |
| Purity | >85% | | |
| Quality Control | The quality of the peptide was evaluated by reversed-phase HPLC and by mass spectrometry. | | |
| Directions for Use | Blocking Peptide to the diluted primary antibody in a molar ratio of 10:1 (peptide to antibody) and incubate the mixture at 4°C for overnight or at room temperature for 2 hours. | | |
| Storage/Stability | Shipped at 4°C. Store at -20°C for one year. | | |

Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC- Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference

Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb- Rabbit, S- Sheep, Z- Zebrafish

COHESION BIOSCIENCES LIMITED

WEB
www.cohesionbio.com

ORDER
order@cohesionbio.com

SUPPORT
techsupport@cohesionbio.com

CUSTOM
custom@cohesionbio.com