

Product Data Sheet

TPL2 Blocking Peptide

Catalog # Source Reactivity Applications

CBP4457 Synthetic H, M, R, B, Mk BL

Description The peptide is used to block Anti-TPL2 Antibody (#CPA4457) reactivity.

Form Lyophilized powder

Gene Symbol MAP3K8

Alternative Names COT; ESTF; Mitogen-activated protein kinase kinase kinase 8; Cancer Osaka thyroid

oncogene; Proto-oncogene c-Cot; Serine/threonine-protein kinase cot; Tumor

progression locus 2; TPL-2

Entrez Gene 1326 (Human); 26410 (Mouse); 116596 (Rat)

SwissProt P41279 (Human); Q07174 (Mouse); Q63562 (Rat)

Purity >85%

Quality Control The quality of the peptide was evaluated by reversed-phase HPLC and by mass

spectrometry.

Directions for UseBlocking Peptide to the diluted primary antibody in a molar ratio of 10:1 (peptide to

antibody) and incubate the mixture at 4°C for overnight or at room temperature for

2 hours.

Storage/Stability Shipped at 4°C. Store at -20°C for one year.

Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC- Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb-Rabbit, S- Sheep, Z- Zebrafish

COHESION BIOSCIENCES LIMITED