

Product Data Sheet

MRPL40 Blocking Peptide

Catalog #	Source	Reactivity	Applications			
CBP4468	Synthetic	Н, М	BL			
Description	The p	eptide is used to bloc	ck Anti-MRPL40 Antibody (#CPA4468) reactivity.			
Form	Lyoph	ilized powder				
Gene Symbol	MRPL	MRPL40				
Alternative Na	ames NLVCF	NLVCF; URIM; 39S ribosomal protein L40, mitochondrial; L40mt; MRP-L40; Nuclear				
	localiz	localization signal-containing protein deleted in velocardiofacial syndrome;				
	Up-re	Up-regulated in metastasis				
Entrez Gene	64976	64976 (Human); 18100 (Mouse)				
SwissProt	Q9NQ	Q9NQ50 (Human); Q9Z2Q5 (Mouse)				
Purity	>85%	>85%				
Quality Contro	ol The qu	The quality of the peptide was evaluated by reversed-phase HPLC and by mass				
	spectr	rometry.				
Directions for	Use Blocki	Blocking Peptide to the diluted primary antibody in a molar ratio of 10:1 (peptide to				
	antibo	ody) and incubate the	e mixture at 4°C for overnight or at room temperature for			
	2 hou	rs.				
Storage/Stabi	lity Shipp	ed at 4°C. Store at -20	0°C for one year.			

Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC-Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb-Rabbit, S- Sheep, Z- Zebrafish

COHESION BIOSCIENCES LIMITED

WEB	ORDER	SUPPORT	CUSTOM
www.cohesionbio.com	order@cohesionbio.com	techsupport@cohesionbio.com	custom@cohesionbio.com