

## EPHA2 (pY588/596) Blocking Peptide

Catalog #	Source	Reactivity	Applications
CBP4632	Synthetic	H, M, R, C, Mk	BL
<b>Description</b>	The peptide is used to block Anti-EPHA2 (pY588/596) Antibody (#CPA4632) reactivity.		
<b>Form</b>	Lyophilized powder		
<b>Gene Symbol</b>	EPHA2; EPHA3; EPHA4		
<b>Alternative Names</b>	EPHA2; ECK; Ephrin type-A receptor 2; Epithelial cell kinase; Tyrosine-protein kinase receptor ECK; EPHA3; ETK; ETK1; HEK; TYRO4; Ephrin type-A receptor 3; EPH-like kinase 4; EK4; hEK4; HEK; Human embryo kinase; Tyrosine-protein kinase TYRO4; Tyrosine-pro		
<b>Entrez Gene</b>	1969; 2042; 2043 (Human); 13836; 13838 (Mouse); 29210 (Rat)		
<b>SwissProt</b>	P29317; P29320; P54764 (Human); Q03145; P29319; Q03137 (Mouse); O08680 (Rat)		
<b>Purity</b>	>85%		
<b>Quality Control</b>	The quality of the peptide was evaluated by reversed-phase HPLC and by mass spectrometry.		
<b>Directions for Use</b>	Blocking Peptide to the diluted primary antibody in a molar ratio of 10:1 (peptide to antibody) and incubate the mixture at 4°C for overnight or at room temperature for 2 hours.		
<b>Storage/Stability</b>	Shipped at 4°C. Store at -20°C for one year.		

**Application key:** E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC- Immunocytochemistry, IP- Immunoprecipitation, CHIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference

**Species reactivity key:** H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb- Rabbit, S- Sheep, Z- Zebrafish

### COHESION BIOSCIENCES LIMITED

**WEB**  
www.cohesionbio.com

**ORDER**  
order@cohesionbio.com

**SUPPORT**  
techsupport@cohesionbio.com

**CUSTOM**  
custom@cohesionbio.com