

**PKA C alpha/beta/gamma (pT198) Blocking Peptide**

Catalog #	Source	Reactivity	Applications
CBP4781	Synthetic	H, M, R, B, D, P, S	BL
<b>Description</b>	The peptide is used to block Anti-PKA C alpha/beta/gamma (pT198) Antibody (#CPA4781) reactivity.		
<b>Form</b>	Lyophilized powder		
<b>Gene Symbol</b>	PRKACA; PRKACB		
<b>Alternative Names</b>	PRKACA; PKACA; cAMP-dependent protein kinase catalytic subunit alpha; PKA C-alpha; PRKACB; cAMP-dependent protein kinase catalytic subunit beta; PKA C-beta; PRKACG; cAMP-dependent protein kinase catalytic subunit gamma; PKA C-gamma		
<b>Entrez Gene</b>	5566; 5567 (Human); 18747; 18749 (Mouse); 293508 (Rat)		
<b>SwissProt</b>	P17612; P22694; P22612 (Human); P05132; P68181 (Mouse); P27791; P68182 (Rat)		
<b>Purity</b>	>85%		
<b>Quality Control</b>	The quality of the peptide was evaluated by reversed-phase HPLC and by mass spectrometry.		
<b>Directions for Use</b>	Blocking Peptide to the diluted primary antibody in a molar ratio of 10:1 (peptide to antibody) and incubate the mixture at 4°C for overnight or at room temperature for 2 hours.		
<b>Storage/Stability</b>	Shipped at 4°C. Store at -20°C for one year.		

**Application key:** E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC- Immunocytochemistry, IP- Immunoprecipitation, CHIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference

**Species reactivity key:** H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb- Rabbit, S- Sheep, Z- Zebrafish

**COHESION BIOSCIENCES LIMITED**

**WEB**  
[www.cohesionbio.com](http://www.cohesionbio.com)

**ORDER**  
[order@cohesionbio.com](mailto:order@cohesionbio.com)

**SUPPORT**  
[techsupport@cohesionbio.com](mailto:techsupport@cohesionbio.com)

**CUSTOM**  
[custom@cohesionbio.com](mailto:custom@cohesionbio.com)