

Product Data Sheet

Dematin Blocking Peptide

| Catalog # | Source | Reactivity | Applications | | | |
|----------------|---------------------|---|---|--|--|--|
| CBP5116 | Synthetic | Н, М, В | BL | | | |
| Description | The p | The peptide is used to block Anti-Dematin Antibody (#CPA5116) reactivity. | | | | |
| Form | Lyop | hilized powder | | | | |
| Gene Symbol | DMT | DMTN | | | | |
| Alternative N | ames DMT | DMT; EPB49; Dematin; Dematin actin-binding protein; Erythrocyte membrane | | | | |
| | prote | ein band 4.9 | | | | |
| Entrez Gene | 2039 | 2039 (Human); 13829 (Mouse) | | | | |
| SwissProt | Q084 | Q08495 (Human); Q9WV69 (Mouse) | | | | |
| Purity | >85% | >85% | | | | |
| Quality Contr | ol The o | The quality of the peptide was evaluated by reversed-phase HPLC and by mass | | | | |
| | spec | trometry. | | | | |
| Directions for | Use Block | Blocking Peptide to the diluted primary antibody in a molar ratio of 10:1 (peptide to | | | | |
| | antib | oody) and incubate the | e mixture at 4°C for overnight or at room temperature for | | | |
| | 2 ho | urs. | | | | |
| Storage/Stab | i lity Shipp | oed at 4°C. Store at -2 | 0°C for one year. | | | |

Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC-Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb-Rabbit, S- Sheep, Z- Zebrafish

COHESION BIOSCIENCES LIMITED

| WEB | ORDER | SUPPORT | CUSTOM |
|---------------------|-----------------------|-----------------------------|------------------------|
| www.cohesionbio.com | order@cohesionbio.com | techsupport@cohesionbio.com | custom@cohesionbio.com |