

Product Data Sheet

HMGN2 (pS29) Blocking Peptide

| Catalog # | Source | Reactivity | Applications | | | |
|-----------------------|-------------|---|---|--|--|--|
| CBP5882 | Synthetic | H, M, R, B, C, D, P | BL | | | |
| Description | The pe | The peptide is used to block Anti-HMGN2 (pS29) Antibody (#CPA5882) reactivity. | | | | |
| Form | Lyoph | ilized powder | | | | |
| Gene Symbol | HMGN | HMGN2 | | | | |
| Alternative Na | ames HMG1 | HMG17; Non-histone chromosomal protein HMG-17; High mobility group | | | | |
| | nucleo | osome-binding domain-c | ontaining protein 2 | | | |
| Entrez Gene | 3151 (| 3151 (Human); 15331 (Mouse) | | | | |
| SwissProt | P0520 | P05204 (Human); P09602 (Mouse); P18437 (Rat) | | | | |
| Purity | >85% | | | | | |
| Quality Contro | ol The qu | The quality of the peptide was evaluated by reversed-phase HPLC and by mass | | | | |
| | spectr | ometry. | | | | |
| Directions for | Use Blocki | Blocking Peptide to the diluted primary antibody in a molar ratio of 10:1 (peptide to | | | | |
| | antibo | dy) and incubate the mi | xture at 4°C for overnight or at room temperature for | | | |
| | 2 hou | ſS. | | | | |
| Storage/Stabi | lity Shippe | ed at 4°C. Store at -20°C | for one year. | | | |

Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC-Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb-Rabbit, S- Sheep, Z- Zebrafish

COHESION BIOSCIENCES LIMITED

| WEB | ORDER | SUPPORT | CUSTOM |
|---------------------|-----------------------|-----------------------------|------------------------|
| www.cohesionbio.com | order@cohesionbio.com | techsupport@cohesionbio.com | custom@cohesionbio.com |