

Product Data Sheet

PER2 (pS662) Blocking Peptide

Catalog #	Source	Reactivity	Applications			
CBP7141	Synthetic	Н, М	BL			
Description	The pe	The peptide is used to block Anti-PER2 (pS662) Antibody (#CPA7141) reactivity.				
Form	Lyoph	ilized powder				
Gene Symbol	PER2	PER2				
Alternative Na	ames KIAAO	KIAA0347; Period circadian protein homolog 2; hPER2; Circadian clock protein				
	PERIO	D 2				
Entrez Gene	8864 (8864 (Human); 18627 (Mouse)				
SwissProt	O1505	55 (Human); O54943	(Mouse)			
Purity	>85%	>85%				
Quality Control	ol The qu	The quality of the peptide was evaluated by reversed-phase HPLC and by mass				
	spectr	ometry.				
Directions for	Use Blocki	Blocking Peptide to the diluted primary antibody in a molar ratio of 10:1 (peptide to				
	antibo	ody) and incubate the	e mixture at 4°C for overnight or at room temperature for			
	2 hou	rs.				
Storage/Stabi	lity Shippe	ed at 4°C. Store at -2	0°C for one year.			

Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC-Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb-Rabbit, S- Sheep, Z- Zebrafish

COHESION BIOSCIENCES LIMITED

WEB	ORDER	SUPPORT	CUSTOM
www.cohesionbio.com	order@cohesionbio.com	techsupport@cohesionbio.com	custom@cohesionbio.com