

# **Product Data Sheet**

# Anti-CD284 Antibody-PE labled

Catalog #	Source	e Reactivity	Applications			
CFF8732	Mouse	e H	IF, FC			
Description		Mouse monoclonal antiboo	ly PE labled to CD284			
Immunogen		Ba/F3 cell line expressing h	uman TLR4			
Purification		The antibody was purified l	by affinity chromatography.			
Specificity		Recognizes human CD284				
Clonality		Monoclonal (clone: HTA12	5)			
Conjugation		PE				
Form		Mouse IgG2a kappa. Liquid	in PBS, pH 7.3, 0.2% BSA, and 0.02% sodium azide.			
Dilution		10 μl / assay				
Gene Symbol		TLR4				
Alternative Names		Toll-like receptor 4; hToll; CD284				
Entrez Gene		7099 (Human)				
SwissProt		O00206 (Human)				
Directions for Use		1. Take 100 $\mu l$ peripheral blood anticoagulated by EDTA and add to the bottom of 5				
		ml tube.				
		2. Add 10 $\mu$ l labeled antibo	dy to the bottom of flow tube mixing with the whole			
		blood, incubate for 20 minu	utes at room temperature away from light.			
		3. Add 2 ml RBC lysis buffer	; incubate for 10 minutes away from light after mixing,			
		dissolve red blood cells.				
		4. Sample tube is set to 100	00 rpm centrifugation for 5 minutes, discard the			
		supernatant.				
		5. Add 2 ml PBS wash buffe	r to resuspend the cells, then 1000 rpm centrifugation			
Application kov		M/P Mastern blot IH Immuno	histochemistry IE- Immunofluorescence, EC- Elow cytometry IC-			

Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC-Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb-Rabbit, S- Sheep, Z- Zebrafish

### **COHESION BIOSCIENCES LIMITED**

WEB	ORDER	SUPPORT	CUSTOM
www.cohesionbio.com	order@cohesionbio.com	techsupport@cohesionbio.com	custom@cohesionbio.com



### **Product Data Sheet**

for 5 minutes, discard the supernatant.

6. Add 0.5 ml PBS wash buffer to resuspend the cells and detect by flow cytometry (sample should be determined on the day on the machine and can also be added fixation overnight at 4 °C then measured).

**Storage/Stability** Shipped and store at 4°C for one year. Do not freeze.

Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC-Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb-Rabbit, S- Sheep, Z- Zebrafish

#### **COHESION BIOSCIENCES LIMITED**

WEBORDERSUPPORTCUSTOMwww.cohesionbio.comorder@cohesionbio.comtechsupport@cohesionbio.comcustom@cohesionbio.com