

## **Product Data Sheet**

# **Anti-CD11a Antibody-PE labled**

Catalog #	Source	Reactivity	Applications
-----------	--------	------------	--------------

CFF8774 Rat M IF, FC

**Description** Rat monoclonal antibody PE labled to CD11a

Immunogen C57BL/6 mouse splenic secondary cytotoxic T cells

**Purification** The antibody was purified by affinity chromatography.

**Specificity** Recognizes mouse CD11a

Clonality Monoclonal (clone: I21/7)

**Conjugation** PE

Form Rat IgG2a kappa. Liquid in PBS, pH 7.3, 0.2% BSA, and 0.02% sodium azide.

Dilution 10 μl / assay

Gene Symbol ITGAL

Alternative Names CD11A; Integrin alpha-L; CD11 antigen-like family member A; Leukocyte adhesion

glycoprotein LFA-1 alpha chain; LFA-1A; Leukocyte function-associated molecule 1

alpha chain; CD11a

Entrez Gene 16408 (Mouse)

SwissProt P24063 (Mouse)

Directions for Use 1. Take 100 μl peripheral blood anticoagulated by EDTA and add to the bottom of 5

ml tube.

2. Add 10  $\mu$ l labeled antibody to the bottom of flow tube mixing with the whole

blood, incubate for 20 minutes at room temperature away from light.

3. Add 2 ml RBC lysis buffer, incubate for 10 minutes away from light after mixing,

dissolve red blood cells.

4. Sample tube is set to 1000 rpm centrifugation for 5 minutes, discard the

Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC-Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference

Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb-Rabbit, S- Sheep, Z- Zebrafish

### **COHESION BIOSCIENCES LIMITED**

WEB ORDER SUPPORT CUSTOM
www.cohesionbio.com order@cohesionbio.com techsupport@cohesionbio.com custom@cohesionbio.com





## **Product Data Sheet**

supernatant.

5. Add 2 ml PBS wash buffer to resuspend the cells, then 1000 rpm centrifugation for 5 minutes, discard the supernatant.

6. Add 0.5 ml PBS wash buffer to resuspend the cells and detect by flow cytometry (sample should be determined on the day on the machine and can also be added fixation overnight at 4 °C then measured).

Storage/Stability

Shipped and store at 4°C for one year. Do not freeze.

Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC-Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb-Rabbit, S- Sheep, Z- Zebrafish

#### **COHESION BIOSCIENCES LIMITED**