

# **Product Data Sheet**

# Anti-CD102 Antibody-PE labled

Catalog #	Source	e Reactivity	Applications			
CFF8841	Rat	Μ	IF, FC			
Description		Rat monoclonal antibody PE	labled to CD102			
Immunogen		COS cells transfected with m	nouse ICAM-2 cDNA			
Purification		The antibody was purified b	y affinity chromatography.			
Specificity		Recognizes mouse CD102				
Clonality		Monoclonal (clone: 3C4)				
Conjugation		PE				
Form		Rat IgG2a kappa. Liquid in PBS, pH 7.3, 0.2% BSA, and 0.02% sodium azide.				
Dilution		10 μl / assay				
Gene Symbol		ICAM2				
Alternative Names		Intercellular adhesion molecule 2; ICAM-2; CD102				
Entrez Gene		15896 (Mouse)				
SwissProt		P35330 (Mouse)				
Directions for Use		1. Take 100 $\mu l$ peripheral blood anticoagulated by EDTA and add to the bottom of 5				
		ml tube.				
		2. Add 10 μl labeled antiboc	ly to the bottom of flow tube mixing with the whole			
		blood, incubate for 20 minu	tes at room temperature away from light.			
		3. Add 2 ml RBC lysis buffer,	incubate for 10 minutes away from light after mixing,			
		dissolve red blood cells.				
		4. Sample tube is set to 100	0 rpm centrifugation for 5 minutes, discard the			
		supernatant.				
		5. Add 2 ml PBS wash buffer	to resuspend the cells, then 1000 rpm centrifugation			
A secold second second second			interchangistry, IC, Interconception of the second se			

Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC-Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb-Rabbit, S- Sheep, Z- Zebrafish

### **COHESION BIOSCIENCES LIMITED**

WEB	ORDER	SUPPORT	CUSTOM
www.cohesionbio.com	order@cohesionbio.com	techsupport@cohesionbio.com	custom@cohesionbio.com



### **Product Data Sheet**

for 5 minutes, discard the supernatant.

6. Add 0.5 ml PBS wash buffer to resuspend the cells and detect by flow cytometry (sample should be determined on the day on the machine and can also be added fixation overnight at 4 °C then measured).

**Storage/Stability** Shipped and store at 4°C for one year. Do not freeze.

Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC-Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb-Rabbit, S- Sheep, Z- Zebrafish

#### **COHESION BIOSCIENCES LIMITED**

WEBORDERSUPPORTCUSTOMwww.cohesionbio.comorder@cohesionbio.comtechsupport@cohesionbio.comcustom@cohesionbio.com