

# **Product Data Sheet**

# Anti-Ly-6A/E Antibody-PE/Cy5 labled

Catalog #	Source	e Reactivity	Applications			
CFG8994	Rat	Μ	IF, FC			
Description	Description Rat monoclonal antibody PE/Cy5 labled to Ly-6A/E					
Immunogen		IL-2 dependent mouse (C57BL/6) cytotoxic T cell line CTL-L				
Purification		The antibody was purified	by affinity chromatography.			
Specificity		Recognizes mouse Ly-6A/E				
Clonality		Monoclonal (clone: D7)				
Conjugation		PE/Cy5				
Form		Rat IgG2a kappa. Liquid in PBS, pH 7.3, and 0.02% sodium azide.				
Dilution		10 μl / assay				
Gene Symbol		Ly6a; Ly6e				
Alternative Names		Ly6; Lymphocyte antigen 6A-2/6E-1; Ly-6A.2/Ly-6E.1; Stem cell antigen 1; SCA-1;				
		T-cell-activating protein; TA	P; Ly67; Sca-2; Tsa-1; Lymphocyte antigen 6E; Ly-6E; Stem			
		cell antigen 2; Thymic share	ed antigen 1; TSA-1			
Entrez Gene		110454, 17069 (Mouse)				
SwissProt		P05533, Q64253 (Mouse)				
<b>Directions for Use</b>		1. Take 100 $\mu l$ peripheral blood anticoagulated by EDTA and add to the bottom of 5				
		ml tube.				
		2. Add 10 µl labeled antibo	dy to the bottom of flow tube mixing with the whole			
		blood, incubate for 20 min	utes at room temperature away from light.			
		3. Add 2 ml RBC lysis buffer	; incubate for 10 minutes away from light after mixing,			
		dissolve red blood cells.				
		4. Sample tube is set to 100	00 rpm centrifugation for 5 minutes, discard the			

Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC-Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb-Rabbit, S- Sheep, Z- Zebrafish

### **COHESION BIOSCIENCES LIMITED**

WEB	ORDER	SUPPORT	CUSTOM
www.cohesionbio.com	order@cohesionbio.com	techsupport@cohesionbio.com	custom@cohesionbio.com



### **Product Data Sheet**

supernatant.

5. Add 2 ml PBS wash buffer to resuspend the cells, then 1000 rpm centrifugation for 5 minutes, discard the supernatant.

6. Add 0.5 ml PBS wash buffer to resuspend the cells and detect by flow cytometry (sample should be determined on the day on the machine and can also be added fixation overnight at 4 °C then measured).

**Storage/Stability** Shipped and store at 4°C for one year. Do not freeze.

Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC-Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb-Rabbit, S- Sheep, Z- Zebrafish

#### **COHESION BIOSCIENCES LIMITED**

WEBORDERSUPPORTCUSTOMwww.cohesionbio.comorder@cohesionbio.comtechsupport@cohesionbio.comcustom@cohesionbio.com