

# **Product Data Sheet**

# Anti-CD86 Antibody-AF700 labled

Catalog #	Source	e Reactivity	Applications		
CFO8834	Rat	М	IF, FC		
Description		Rat monoclonal antibody Al	700 labled to CD86		
Immunogen		LPS-activated CBA/Ca mous	e splenic B cells		
Purification		The antibody was purified b	y affinity chromatography.		
Specificity		Recognizes mouse CD86			
Clonality		Monoclonal (clone: GL-1)			
Conjugation		AF700			
Form		Rat IgG2a kappa. Liquid in PBS, pH 7.3, 0.2% BSA, and 0.02% sodium azide.			
Dilution		10 μl / assay			
Gene Symbol		CD86			
Alternative Names		CD28LG2; T-lymphocyte activation antigen CD86; Activation B7-2 antigen; B70;			
		BU63; CTLA-4 counter-recep	otor B7.2; FUN-1; CD86		
Entrez Gene		12524 (Mouse)			
SwissProt		P42082 (Mouse)			
Directions for Use		1. Take 100 $\mu l$ peripheral blood anticoagulated by EDTA and add to the bottom of 5			
		ml tube.			
		2. Add 10 $\mu$ l labeled antibod	ly to the bottom of flow tube mixing with the whole		
		blood, incubate for 20 minu	tes at room temperature away from light.		
		3. Add 2 ml RBC lysis buffer,	incubate for 10 minutes away from light after mixing,		
		dissolve red blood cells.			
		4. Sample tube is set to 100	0 rpm centrifugation for 5 minutes, discard the		
		supernatant.			

Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC-Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb-Rabbit, S- Sheep, Z- Zebrafish

### **COHESION BIOSCIENCES LIMITED**

WEB	ORDER	SUPPORT	CUSTOM
www.cohesionbio.com	order@cohesionbio.com	techsupport@cohesionbio.com	custom@cohesionbio.com



### **Product Data Sheet**

5. Add 2 ml PBS wash buffer to resuspend the cells, then 1000 rpm centrifugation for 5 minutes, discard the supernatant.

6. Add 0.5 ml PBS wash buffer to resuspend the cells and detect by flow cytometry (sample should be determined on the day on the machine and can also be added fixation overnight at 4 °C then measured).

**Storage/Stability** Shipped and store at 4°C for one year. Do not freeze.

Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC-Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb-Rabbit, S- Sheep, Z- Zebrafish

#### **COHESION BIOSCIENCES LIMITED**

WEBORDERSUPPORTCUSTOMwww.cohesionbio.comorder@cohesionbio.comtechsupport@cohesionbio.comcustom@cohesionbio.com