

Product Data Sheet

Anti-CD38 Antibody-Pacific Blue labled

Catalog #	Source	Reactivity	Applications		
CFP8523	Mouse	Н	IF, FC		
Description Mouse monoclonal antibody Pacific Blue labled to CD38					
Immunogen		Native purified human CD3	8.		
Purification		The antibody was purified by affinity chromatography.			
Specificity		Recognizes human CD38			
Clonality		Monoclonal (clone: HIT2)			
Conjugation		Pacific Blue			
Form Mouse IgG1. Liquid in PBS, pH 7.3, 0.2% BSA, and 0.02% sodium azide.					
Dilution 10 μl / assay					
Gene Symbol		CD38			
Alternative Names		ADP-ribosyl cyclase/cyclic ADP-ribose hydrolase 1; 2'-phospho-ADP-ribosyl cyclase;			
		2'-phospho-ADP-ribosyl cyo	clase/2'-phospho-cyclic-ADP-ribose transferase;		
		2'-phospho-cyclic-ADP-ribc	se transferase; ADP-ribosyl cyclase 1; ADPRC 1; Cyclic		
		ADP-ribose hydrola			
Entrez Gene		952 (Human)			
SwissProt	rot P28907 (Human)				
Directions for	Use	1. Take 100 μl peripheral blood anticoagulated by EDTA and add to the bottom of 5			
		ml tube.			
		2. Add 10 μl labeled antibo	dy to the bottom of flow tube mixing with the whole		
		blood, incubate for 20 min	utes at room temperature away from light.		
		3. Add 2 ml RBC lysis buffe	; incubate for 10 minutes away from light after mixing,		
		dissolve red blood cells.			

Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC-Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb-Rabbit, S- Sheep, Z- Zebrafish

COHESION BIOSCIENCES LIMITED

WEB	ORDER	SUPPORT	CUSTOM
www.cohesionbio.com	order@cohesionbio.com	techsupport@cohesionbio.com	custom@cohesionbio.com



Product Data Sheet

4. Sample tube is set to 1000 rpm centrifugation for 5 minutes, discard the supernatant.

5. Add 2 ml PBS wash buffer to resuspend the cells, then 1000 rpm centrifugation for 5 minutes, discard the supernatant.

6. Add 0.5 ml PBS wash buffer to resuspend the cells and detect by flow cytometry (sample should be determined on the day on the machine and can also be added fixation overnight at 4 °C then measured).

Storage/Stability Shipped and store at 4°C for one year. Do not freeze.

Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC-Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb-Rabbit, S- Sheep, Z- Zebrafish

COHESION BIOSCIENCES LIMITED

WEBORDERSUPPORTCUSTOMwww.cohesionbio.comorder@cohesionbio.comtechsupport@cohesionbio.comcustom@cohesionbio.com