

Product Data Sheet

Prestained Protein Ladder (10-250 kDa)

Catalog #	Source	e Reactivity	Applications
CRG1032			WB, Protein Electrophoresis
Description		Prestained Protein Ladder	10 to 250 kDa
Specificity		Prestained Protein Ladder	s a mixture of 11 blue-, orange-, green-stained proteins
		(10 to 250 kDa) for use as s	ize standards in protein electrophoresis (SDS-PAGE) and
		western blotting. The prote	ein ladder is supplied in a ready-to-use format for direct
		loading onto gels; no need	to heat, reduce, or add sample buffer prior to use.
Form		Liquid in 67 mM Tris-H3PO	4, 5mM EDTA, 2 % SDS, 33 % Glycerol, 0.02 % proclin300,
		pH7.5	
Directions for Use		1. Thaw the ladder at room temperature for a few minutes to dissolve precipitated	
		solids. Do not boil!	
		2. Mix gently, but thorough	ly, to ensure the solution is homogeneous.
		3. Load the following volun	nes of the ladder on an SDS-polyacrylamide gel:
		- 5 μ L per well for mini gel,	
		- 10 μL per well for large ge	l.
		Use the same volumes for	Western blotting.
		The loading volumes listed	above are recommended for gels with a thickness of

Notes:

1. Prestained proteins can have different mobilities in various SDS-PAGE-buffer systems. However, they are suitable for approximate molecular weight determination when calibrated against unstained standards in the same system. See the table provided for migration patterns in different electrophoresis conditions. In low-percentage gels (< 10 %), the low-molecular weight proteins in the ladder may migrate with the dye front.

0.75-1.0 mm. The loading volume should be doubled for 1.5 mm thick gels.

2. Prestained Protein Ladder can be used in Western blotting with all common

COHESION BIOSCIENCES LIMITED



Product Data Sheet

membranes: PVDF, nylon and nitrocellulose.

3. Longer transfer times or higher transfer voltages may be required for Western blotting of large (>100 kDa) proteins.

Storage/Stability

Image

Shipped at 4°C. Store at -20°C for one year.

