

Product Data Sheet

FDX1 siRNA (Mouse)

Catalog #	Source	Reactivity	Applications		
CRM1331	Synthetic	M	RNAi		
	-				
Description	siRNA	siRNA to inhibit FDX1 expression using RNA interference			
Specificity	FDX1	FDX1 siRNA (Mouse) is a target-specific 19-23 nt siRNA oligo duplexes designed to			
kno		knock down gene expression.			
Form	Lyoph	Lyophilized powder			
Gene Symbol FDX		FDX1			
Alternative Na	tive Names Adrenodoxin mitochondrial; Adrenal ferredoxin; Ferredoxin-1			1	
Entrez Gene	Entrez Gene 14148 (Mouse)				
SwissProt P46656 (Mouse)					
Purity > 97%					
Quality Contro	Oligonucleotide synthesis is monitored base by base through trityl analysis to e			n trityl analysis to ensure	
	appro	appropriate coupling efficiency. The oligo is subsequently purified by affinity-so			
phase extraction. The annealed RNA duplex is further analyzed by mass			ed by mass		
	specti	spectrometry to verify the exact composition of the duplex. Each lot is compared to			
	the pr	revious lot by mass sp	ectrometry to ensure maximum l	ot-to-lot consistency.	
Components	We of	fers pre-designed sets	rs pre-designed sets of 3 different target-specific siRNA oligo duplexes of		
mouse FDX1 gene. Each vial contains 5 nmol of lyophilized siRNA. The c be transfected individually or pooled together to achieve knockdown o			l contains 5 nmol of lyophilized si	RNA. The duplexes can	
			ockdown of the target		
	gene, which is most commonly assessed by qPCR or western blot.			blot.	
Component 15			15 nmol	30 nmol	
	FDX1	L siRNA (Mouse) - A	5 nmol x 1	5 nmol x 2	

Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC-Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb-Rabbit, S- Sheep, Z- Zebrafish

5 nmol x 1

5 nmol x 2

FDX1 siRNA (Mouse) - B

COHESION BIOSCIENCES LIMITED

WEB	ORDER	SUPPORT	CUSTOM
www.cohesionbio.com	order@cohesionbio.com	techsupport@cohesionbio.com	custom@cohesionbio.com



Product Data Sheet

-			
	DEPC Water	1 ml x 1	1 ml x 2
	Negative Control	2.5 nmol x 1	2.5 nmol x 2
	FDX1 siRNA (Mouse) - C	5 nmol x 1	5 nmol x 2

Directions for Use

We recommends transfection with 10 nM - 100 nM siRNA 48 to 72 hours prior to cell lysis. Before resuspending, briefly centrifuge the tube to ensure the lyophilized siRNA is at the bottom of the tube. Resuspend the siRNA oligos to an appropriate concentration with DEPC water. For example, resuspend one tube of 5 nmol siRNA oligo in 250 μ l of DEPC water to get a final concentration of 20 μ M.

Plate	Final volume	Final concentration	siRNA (20 μM)	Lipofectamin
	of medium	of siRNA		2000
		100 nM	0.5 μl	0.25 μl
96-well	100 µl	50 nM	0.25 μl	0.25 μl
		10 nM	0.05 μl	0.25 μl
		100 nM	2.5 μl	1 µl
24-well	500 μl	50 nM	1.25 μl	1 µl
		10 nM	0.25 μl	1 µl
		100 nM	5 μl	2 µl
12-well	1 ml	50 nM	2.5 μl	2 µl
		10 nM	0.5 μl	2 µl
		100 nM	10 µl	5 µl
6-well	2 ml	50 nM	5 μl	5 µl
		10 nM	1 µl	5 µl

Storage/Stability

Shipped at 4 °C. Store at -20 °C for one year.

Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC-Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb-Rabbit, S- Sheep, Z- Zebrafish

COHESION BIOSCIENCES LIMITED

WEB	ORDER	SUPPORT	CUSTOM
www.cohesionbio.com	order@cohesionbio.com	techsupport@cohesionbio.com	custom@cohesionbio.com