

Storage/Stability

Product Data Sheet

Rabbit Anti-Dog IgG (H&L) - APC

freeze/thaw cycles.

Catalog #	Source	Reactivity	Applications	
CSA4815	Rabbit	D	E, IH, IF, FC	
Description	Rab	Rabbit Polyclonal Secondary Antibody to Dog IgG (H&L) APC labeled		
Immunogen	Dog	Dog IgG		
Purification	The	The antibody was isolated from antisera by immunoaffinity chromatography using		
	antigens coupled to agarose beads.			
Specificity	Ву і	By immunoelectrophoresis and ELISA this antibody reacts specifically with Dog IgG.		
	No	antibody was detected	against non immunoglobulin serum proteins.	
Clonality	Poly	Polyclonal		
Conjugation	APC	APC		
Form	1 m	ıg/ml. Liquid in 0.01M F	Phosphate Buffered Saline, pH 7.2, containing 1% BSA,	
	50%	6 glycerol, 0.02% Sodiui	m Azide	
Dilution	IH (1/100 - 1/1000), IF (1/100 - 1/1000), FC (1/1000 - 1/4000), E (Use at an assay			
	dep	endent concentration)		
Fluorophore	Ex/	Em = 651/662 nm		

Shipped at 4°C. Upon delivery aliquot and store at -20°C for one year. Avoid

Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC-Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb-Rabbit, S- Sheep, Z- Zebrafish

COHESION BIOSCIENCES LIMITED

WEB ORDER SUPPORT CUSTOM
www.cohesionbio.com order@cohesionbio.com techsupport@cohesionbio.com custom@cohesionbio.com